

# Solution to Assignment 10

Com S 477/577


Nov 19, 2010

1. (courtesy Wen-Chieh Chang)

Please see the Assignments page for an example.

The function has a local minimum  $-3.46410 = -2\sqrt{3}$  at  $(\sqrt{3}, 0)$  and a local maximum  $3.46410 = 2\sqrt{3}$  at  $(-\sqrt{3}, 0)$ .

Note that the function  $f(x_1, x_2)$  is a cubic polynomial on  $x_1$  and  $x_2$ , hence the global maximum and global minimum are  $\infty$  and  $-\infty$ . In other words, the start point of the search is important.



2. Apply the method of Lagrange multipliers. We find the Lagrangian and its gradient:

$$\begin{aligned} F(x_1, x_2, x_3, \lambda) &= f(x_1, x_2, x_3) + \lambda h(x_1, x_2, x_3) \\ &= (3x_1^2 + 4x_2^2 + 5x_3^2) + \lambda(x_1 + x_2 + x_3 - 10), \\ \nabla F &= \begin{pmatrix} 6x_1 + \lambda \\ 8x_2 + \lambda \\ 10x_3 + \lambda \\ x_1 + x_2 + x_3 - 10 \end{pmatrix} = 0. \end{aligned}$$

Solving the above system yields the optimal solution  $(x_1, x_2, x_3) = (\frac{200}{47}, \frac{150}{47}, \frac{120}{47})$  while  $\lambda = -\frac{1200}{47}$ .

3. An inscribed parallelepiped has its eight vertices on the ellipsoid. Let  $(x, y, z)$  denote one of these vertices, where  $x, y, z > 0$ . The volume of the parallelepiped is then  $8xyz$ . There fore we are to solve the following constrained optimization problem:

$$\begin{aligned} & \text{maximize} && 8xyz \\ & \text{subject to} && \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1. \end{aligned}$$

At a maximum point, the Lagrangian

$$F(x, y, z, \lambda) = 8xyz + \lambda \left( \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} - 1 \right)$$

must have vanishing partial derivatives with respect to  $x, y, z$ , and to the Lagrange multiplier  $\lambda$  (the first order necessary condition). This yields

$$\frac{\partial F}{\partial x} = 8yz + \lambda \frac{2x}{a^2} = 0; \tag{1}$$

$$\frac{\partial F}{\partial y} = 8xz + \lambda \frac{2y}{b^2} = 0; \tag{2}$$

$$\frac{\partial F}{\partial z} = 8xy + \lambda \frac{2z}{c^2} = 0. \tag{3}$$

Multiply (1) by  $x$  and (2) by  $y$ , and subtract the second resulting equation from the first one. This yields

$$\lambda \left( \frac{2x^2}{a^2} - \frac{2y^2}{b^2} \right) = 0. \tag{4}$$

Similar steps of manipulation applied to the other two combinations of equations (1), (2), (3) yield

$$\lambda \left( \frac{2x^2}{a^2} - \frac{2y^2}{b^2} \right) = 0, \tag{5}$$

$$\lambda \left( \frac{2x^2}{a^2} - \frac{2y^2}{b^2} \right) = 0. \tag{6}$$

Equations (4)-(6) together imply that either  $\lambda = 0$  or

$$\frac{x^2}{a^2} = \frac{y^2}{b^2} = \frac{z^2}{c^2}. \tag{7}$$

Obviously  $\lambda \neq 0$  as otherwise at least two of  $x, y, z$  would have to be zero. Substituting equation (7) back into the equation of the ellipsoid, we obtain

$$\begin{aligned} x &= \frac{a}{\sqrt{3}}; \\ y &= \frac{b}{\sqrt{3}}; \\ z &= \frac{c}{\sqrt{3}}. \end{aligned}$$

Hence the maximum volume is  $\frac{8abc}{3\sqrt{3}}$ , which is achieved at points  $(\pm \frac{a}{\sqrt{3}}, \pm \frac{b}{\sqrt{3}}, \pm \frac{c}{\sqrt{3}})$ .